


Introductory Chemistry
Chem 103

Chapter 4 – Light and Electronic Structure

Lecture Slides



The Electromagnetic Spectrum




Photo credits (clockwise from top left): Anne, Oni/Deposit Photos; Gerald D. Tang / iStockphoto; Stock, Kevin Revell


What is Light?

electromagnetic radiation


- a form of energy
- travels in waves
- exists in increments called **photons**



The Electromagnetic Spectrum



Wavelength



Describing Electromagnetic Waves

wavelength (λ) – The length of one wave

frequency (ν) – The number of waves per second

$$1 \text{ wave/second} = 1 \text{ hertz (Hz)}$$

10,000 Hz

10,000/s

10,000 s⁻¹

Describing Electromagnetic Waves, Continued

↓ wavelength
↑ frequency } *inversely related*

$$c = \lambda \nu$$

speed of light = wavelength x frequency

$$\frac{m}{s} = m \times \frac{1}{s}$$

c = speed of light = 3.00×10^8 m/s

Example of Describing Electromagnetic Waves

A beam of green light has a wavelength of 500 nm.

What is the frequency of this light?

$$c = \lambda \nu$$

$$c = 3.00 \times 10^8 \text{ m/s}$$

$$\lambda = 500 \text{ nm} = 500 \times 10^{-9} \text{ m}$$

$$1 \text{ nm} = 10^{-9} \text{ m}$$

$$\nu = ?$$

$$\frac{c}{\lambda} = \nu$$

$$\frac{3.00 \times 10^8 \text{ m/s}}{500 \times 10^{-9} \text{ m}} = \nu$$

units: 1/s = Hz


$$6 \times 10^{14} \text{ Hz} = \nu$$

Frequency and Wavelength

The energy of light depends on its frequency and wavelength.



longer wavelength
lower frequency
lower energy



shorter wavelength
higher frequency
higher energy

Energy of a photon:



$$E = h\nu$$

energy

frequency

Planck's constant
= $6.63 \times 10^{-34} \text{ J}\cdot\text{s}$

$$\nu = c/\lambda$$

$$E = hc/\lambda$$

Example of Photon Energy

A photon has a frequency of $7.50 \times 10^{14} \text{ Hz}$. What is the wavelength of this light? What color is this light? What is the energy of the photon?


$$c = \lambda \nu$$

$$\frac{c}{\nu} = \lambda$$

$$\frac{3.00 \times 10^8 \text{ m/s}}{7.50 \times 10^{14} / \text{s}} = \lambda$$

$$4.00 \times 10^{-7} \text{ m} = \lambda$$

$$= 400 \text{ nm} \text{ violet}$$




$$E = h\nu$$


$$E = (6.63 \times 10^{-34} \text{ J}\cdot\text{s})(7.50 \times 10^{14} / \text{s})$$

$$E = 4.97 \times 10^{-19} \text{ J}$$

Summary of Electromagnetic Waves



- Light is a form of electromagnetic radiation
- We describe light by its
 - frequency (ν)
 - wavelength (λ)
 - energy (E)
- $c = \lambda \nu$
- $E = h\nu = hc/\lambda$




Color, Line Spectra, and the Bohr Model



Flame Tests


observe colors emitted by different metal ions




Gas lamps also produce unique colors:



Line Spectra




Line Spectra, Continued



Examples of Line Spectra


Each element produces a unique line spectrum.



Photoelectric Effect

Early 20th Century:


- Dense nucleus surrounded by electrons
- *Photoelectric effect*: light causes atoms to eject electrons



Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

The Bohr Model (1913)

- Electrons orbit the nucleus.
- Only certain orbit energies are "allowed".
- Electrons can jump between levels.
- Light is absorbed or released when electrons jump.
- *Ground state*: all electrons in lowest possible levels.




Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

The Bohr Model, Continued




Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company



Revel, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Bohr Model and Line Spectra



The Hydrogen Atom





TABLE 4.1 Transition in the Hydrogen Line Spectrum

Transition	Color Produced
3 → 2	Red
4 → 2	Light blue
5 → 2	Indigo (deep blue)
6 → 2	Purple(violet)

Light and Electrons



Sources of Light



Do You Give Off Light?




Summary of the Bohr Model

Explained

- The hydrogen line spectrum
- Some properties of main group elements


Did not explain

- More complex line spectra
- Properties of the transition elements




The Quantum Model and Electron Orbitals

Bohr Model: 1913



Quantum Model: 1920s-30s



Heisenberg's Uncertainty Principle

It is impossible to precisely know the exact velocity and location of a particle.

We describe the **shape** the blades occupy.




Quantum mechanics: describes electrons { most probable locations
energies

The wave nature of electrons

Tiny, fast-moving particles also behave as waves.

This explains electron energy levels.




The Quantum Model

Main Ideas:

- uncertainty principle
- wave nature of electrons

QM describes electrons by


- energy
- probable locations



Energy Levels and Sublevels, Part 1

1. Electrons occupy different energy levels.

- Level is identified by its **principal quantum number**, n (1, 2, 3...)
- Higher energy levels can hold more electrons



Level	Electron Capacity
1	2
2	8
3	18
4	32

Energy Levels and Sublevels, Part 2

2. Each energy level contains one or more **sublevels**.


Sublevel
s
p
d
f

Energy Levels and Sublevels, Part 3


3. Each sublevel contains one or more **orbitals**.

Sublevel	Number of Orbitals
s	1
p	3
d	5
f	7

The Bohr Model and the Quantum Model



Bohr model
Electrons orbit like planets



Quantum model
Electrons behave like waves that occupy different regions

Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Energy Levels and Sublevels, Part 4

4. Each orbital holds up to two electrons.

- Electrons have a magnetic field, called spin.
- Electrons with opposite spins pair together.




Revell, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Energy Levels and Sublevels, Summary


1. Electrons occupy different energy levels.
2. Each level contains sublevels.
3. Each sublevel contains orbitals.
4. Each orbital holds up to two electrons.

Sublevel	Number of Orbitals	Electron Capacity
<i>s</i>	1	2
<i>p</i>	3	6
<i>d</i>	5	10
<i>f</i>	7	14


Level 1: *s* only



Level 2: *s* + *p*, part 1



Level 2: *s* + *p*, part 2




Level 2: *s* + *p*, part 3

Sublevel	Number of Orbitals	Electron Capacity
<i>s</i>	1	2
<i>p</i>	3	6

Total: 8

Level 3: *s* + *p* + *d*, part 1




Level 3: $s + p + d$, part 2

Sublevel	Number of Orbitals	Electron Capacity
s	1	2
p	3	6
d	5	10

Total: 18

Level 4: $s + p + d + f$, part 1



Level 4: $s + p + d + f$, part 2

Sublevel	Number of Orbitals	Electron Capacity
s	1	2
p	3	6
d	5	10
f	7	14

Total: 32


Summary of Atomic Energy Levels

TABLE 4.4 Energy Levels, Sublevels, and Electron Capacity

Energy Level	1	2	3	4
Sublevels			$d (10 e^-)$	$f (14 e^-)$
		$p (6 e^-)$	$p (6 e^-)$	$p (6 e^-)$
	$s (2 e^-)$	$s (2 e^-)$	$s (2 e^-)$	$s (2 e^-)$
Electron Capacity	2	8	18	32


Note : the symbol e^- means electron.







Energy Differences Between Levels



Describing Electron Configuration


Quantum Model:
Energy levels – 1, 2, 3...
Energy sublevels – s, p, d, f





Hund's Rule:
If empty orbitals of the same energy are available, electrons singly occupy orbitals rather than pairing together.

Carbon:



Electron Configurations of Row 2 Elements

3	4
Li 6.94	Be 9.01


Li: $1s^2 2s^1$
Be: $1s^2 2s^2$

5	6	7	8	9	10
B 10.81	C 12.01	N 14.01	O 16.00	F 19.00	Ne 20.18


B: $1s^2 2s^2 2p^1$
C: $1s^2 2s^2 2p^2$
N: $1s^2 2s^2 2p^3$
O: $1s^2 2s^2 2p^4$
F: $1s^2 2s^2 2p^5$
Ne: $1s^2 2s^2 2p^6$

Example for Silicon

What is the electron configuration of silicon?




Energy Diagram and Writing Electron Configurations



Describing Electron Configuration, Part 2

valence level: The highest occupied electron energy level

- Up to 8 electrons in valence level




Describing Electron Configurations, Part 3

Argon: (18 e⁻)

$1s^2 2s^2 2p^6 3s^2 3p^6$
valence level

Potassium: (19 e⁻)

$1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$
valence level



Noble Gases have Filled Valences

2	He	4.00
10	Ne	20.18
18	Ar	39.95
36	Kr	83.80

$1s^2$

$1s^2 2s^2 2p^6$

$1s^2 2s^2 2p^6 3s^2 3p^6$

$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6$

Octet Rule:


An atom is stabilized by having its highest-occupied (valence) energy level filled.

Electron Configurations for Larger Atoms

	inner electrons	Noble gas notation
Sodium:	$1s^2 2s^2 2p^6 3s^1$	$[\text{Ne}]3s^1$
Phosphorous:	$1s^2 2s^2 2p^6 3s^2 3p^3$	$[\text{Ne}]3s^2 3p^3$
Chlorine:	$1s^2 2s^2 2p^6 3s^2 3p^5$	$[\text{Ne}]3s^2 3p^5$
	$1s^2 2s^2 2p^6 = [\text{Ne}]$	

Electron Configurations for Larger Atoms, Continued



not involved with bonding



Example of Writing an Electron Configuration

Write the electron configuration for selenium using the noble gas shorthand. Identify the inner electrons, the outer electrons, and the valence electrons.

34	Se	78.97
----	----	-------



Example, Electron Configuration for Ions - Sodium

11	Na	22.99
----	----	-------

What is the electron configuration of a sodium atom?

What is the electron configuration of a sodium ion with a +1 charge?

species	Symbol	full configuration	noble-gas shorthand
sodium atom	Na	$1s^2 2s^2 2p^6 3s^1$	$[\text{Ne}]3s^1$
sodium ion (+1 charge)	Na^+	$1s^2 2s^2 2p^6$	$[\text{He}]2s^2 2p^6$ or $[\text{Ne}]$

Example, Electron Configuration for Ions - Oxygen

8	O	16.00
---	---	-------

What is the electron configuration of an oxide ion, which is an oxygen ion with a charge of -2?

species	symbol	full configuration	noble-gas shorthand
oxygen atom	O	$1s^2 2s^2 2p^4$	$[\text{He}]2s^2 2p^4$
oxide ion (-2 charge)	O^{2-}	$1s^2 2s^2 2p^6$	$[\text{He}]2s^2 2p^6$ or $[\text{Ne}]$

Many ions form noble gas configurations

O: $1s^2 2s^2 2p^4$ Na: $1s^2 2s^2 2p^6 3s^1$

O^{2-} : $1s^2 2s^2 2p^6$ Na^+ : $1s^2 2s^2 2p^6$

Ne: $1s^2 2s^2 2p^6$

These are isoelectronic

Electron Configuration and the Periodic Table



Group 1A Electron Configurations



3 Li 6.94 [He]2s ¹
11 Na 22.99 [Ne]3s ¹
19 K 39.10 [Ar]4s ¹

Lithium (3 electrons): [He]2s¹

Sodium (11 electrons): [Ne]3s¹

Potassium (19 electrons): [Ar]4s¹

Group 7A Electron Configurations

Fluorine: [He]2s²2p⁵

Chlorine: [Ne]3s²3p⁵

Bromine: [Ar]4s²3d¹⁰4p⁵

9 F 19.00 [He]2s ² 2p ⁵
17 Cl 35.45 [Ne]3s ² 3p ⁵
35 Br 79.90 [Ar]4s ² 3d ¹⁰ 4p ⁵

Row and Energy Level

1	H	He																
2	Li	Be	B	C	N	O	F	Ne										
3	Na	Mg	Al	Si	P	S	Cl	Ar										
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
6	Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
7	Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Cn	Nh	Fl	Mc	Lv	Ts	Og	

The row indicates the highest occupied electron energy level.

56	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Fm	Tm	Yb
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr


Column and Electron Configuration

	s^1																	p^6					
1	H																	He					
2	Li	Be	s^2															B	C	N	O	F	Ne
3	Na	Mg		d^1	d^2	d^3	d^4	d^5	d^6	d^7	d^8	d^9	d^{10}	Al	Si	P	S	Cl	Ar				
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr					
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe					
6	Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn					
7	Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Cn	Nh	Fl	Mc	Lv	Ts	Og						


The column gives the outermost electron configuration.

	p^1	p^2	p^3	p^4	p^5	p^6													p^1	p^2	p^3	p^4	p^5	p^6
1	B	C	N	O	F	Ne													B	C	N	O	F	Ne
2	Al	Si	P	S	Cl	Ar													Al	Si	P	S	Cl	Ar
3	Ga	Ge	As	Se	Br	Kr													Ga	Ge	As	Se	Br	Kr
4	In	Sn	Sb	Te	I	Xe													In	Sn	Sb	Te	I	Xe
5	Tl	Pb	Bi	Po	At	Rn													Tl	Pb	Bi	Po	At	Rn
6	Pb	Bi	Po	At	Rn														Pb	Bi	Po	At	Rn	
7	Fr	Ra	Ac																Fr	Ra	Ac			

Four Blocks of the Periodic Table




Organization of the Periodic Table



Sulfur Electron Configuration

What is the outermost electron configuration for sulfur?


$3p^4$



Highest-Energy Occupied Sublevel

Write the configuration for the highest-energy occupied sublevel for potassium, phosphorus, and iron.

K: $4s^1$
P: $3p^3$
Fe: $3d^6$




Electron Configuration of Aluminum


Write the electron configuration for aluminum.

How many valence electrons does aluminum have?

$[Ne]3s^23p^1$
3 valence electrons



Valence Electrons



Summary of Periodic Table Organization


The row indicates the highest occupied electron energy level.

The column gives the outermost electron configuration.

Introductory Chemistry Chem 103

Chapter 5 – Chemical Bonds and Compounds

Lecture Slides



Lewis Symbols and the Octet Rule

Valence electrons

- electrons in highest occupied energy level
- s and p sublevels
- generally up to 8 electrons

Lewis Symbols Show Valence Electrons

Lewis dot symbols

- Represent valence electrons as dots around atomic symbol

Group	1A	2A	3A	4A	5A	6A	7A	8A
Valence electrons	1	2	3	4	5	6	7	8
Configuration	s^1	s^2	s^2p^1	s^2p^2	s^2p^3	s^2p^4	s^2p^5	s^2p^6

The Octet Rule

Octet Rule: An atom is stabilized by having its valence energy level filled.



Noble gases fulfill the octet rule.

Other atoms fulfill the octet rule by:


- gaining or losing electrons (ions).
- sharing electrons.

Ions

Atoms or groups of atoms that have an overall charge.



Cations – positively charged ions

Main group metals fulfill the octet rule by forming cations



Lewis Structures Show Cation Formation

+1	+2
H ⁺	Be ²⁺
Li ⁺	Mg ²⁺
Na ⁺	Ca ²⁺
K ⁺	Sr ²⁺
Rb ⁺	



Transition metals also form cations.

Typical charges are +1, +2, +3, or +4

Some metals form multiple charged ions.

p-block metals also do this.

Revised, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Naming Cations

Metal cations have the same name as the neutral metal.

Na⁺ sodium

Mg²⁺ magnesium

Atom	Ion	Older Name	Modern Name
Iron	Fe ²⁺	ferrous	iron(II)
	Fe ³⁺	ferric	iron(III)
Copper	Cu ⁺	cuprous	copper(I)
	Cu ²⁺	cupric	copper(II)

Practice Naming Cations

Name the following cations:

Ag⁺ Pb²⁺ Pb⁴⁺
silver *lead(II)* *lead(IV)*


Revised, Introductory Chemistry, 2e, © 2021 W. H. Freeman and Company

Anions – negatively charged ions

N	O	F	
P	S	Cl	
	Se	Br	
		I	


Anions Fulfill the Octet Rule, Part 1

Most nonmetals fulfill the octet rule by gaining electrons.



Anions Fulfill the Octet Rule, Part 2

Most nonmetals fulfill the octet rule by gaining electrons.



Naming Anions: change ending to *-ide*

Atom	Anion Symbol	Anion Name
chlorine	Cl^-	chloride
oxygen	O^{2-}	oxide
sulfur	S^{2-}	sulfide
nitrogen	N^{3-}	nitride

Polyatomic ions: groups of atoms with a charge, part 1

NH_4^+ Ammonium			
NO_3^-	Nitrate	SO_4^{2-}	Sulfate
NO_2^-	Nitrite	SO_3^{2-}	Sulfite
CO_3^{2-}	Carbonate	HSO_4^-	Bisulfate (Hydrogen sulfate)
HCO_3^-	Bicarbonate (Hydrogen carbonate)	ClO_4^-	Perchlorate
PO_4^{3-}	Phosphate	ClO_3^-	Chlorate
HPO_4^{2-}	Hydrogen phosphate	ClO_2^-	Chlorite
$\text{C}_2\text{H}_3\text{O}_2^-$	Acetate	ClO^-	Hypochlorite
OH^-	Hydroxide	CrO_4^{2-}	Chromate
CN^-	Cyanide	$\text{Cr}_2\text{O}_7^{2-}$	Dichromate
O_2^{2-}	Peroxide	MnO_4^-	Permanganate

Polyatomic ions: groups of atoms with a charge, part 2

Oxyanions – contain oxygen

Usually named as element root + *-ate*

CO_3^{2-} carbonate

PO_4^{3-} phosphate

Polyatomic ions: groups of atoms with a charge, part 3

More than one oxyanion:

-ate more oxygen atoms
-ite fewer oxygen atoms

NO_3^- nitrate

NO_2^- nitrite

Polyatomic ions: groups of atoms with a charge, part 4

More than one oxyanion:

- ate* more oxygen atoms
- ite* fewer oxygen atoms

ClO_4^- perchlorate

ClO_3^- chlorate

ClO_2^- chlorite

ClO^- hypochlorite

-ite fewer oxygen atoms



Ions to Know


Monatomic anions

H ⁺																
Li ⁺	Ba ²⁺															
Na ⁺	Mg ²⁺															
K ⁺	Ca ²⁺															
Rb ⁺	Sr ²⁺															
		Co^{2+} Ni^{2+} Fe^{2+} Cu^{2+} Cr^{3+} Mn^{2+} Fe^{3+} Co^{3+}														
												Zn^{2+}				
												Ag^{+}				
												Au^{3+} Au^{+} Pt^{2+} Pt^{4+}				
												Hg^{2+}				
												Pb^{2+}				
												Bi^{3+}				
												Sb^{3+}				
												Sn^{2+}				
												Sn^{4+}				
												Pb^{4+}				
												Bi^{5+}				
												Sb^{5+}				
												As^{3+}				
												As^{5+}				
												V^{3+}				
												V^{5+}				
												Cr^{6+}				
												Mn^{7+}				
												N^{3-}				
												O^{2-}				
												F^{-}				
												Cl^{-}				
												Br^{-}				
												I^{-}				

Polytatomic anions

NH_4^{+} Ammonium	
NO_3^{-} Nitrate	SO_4^{2-} Sulfate
CO_3^{2-} Carbonate	SO_3^{2-} Sulfite
HCO_3^{-} Bicarbonate	HSO_4^{-} Bisulfate
(Hydrogen carbonate)	(Hydrogen sulfate)
NO_2^{-} Nitrite	CO_3^{2-} Percarbonate
PO_4^{3-} Phosphate	O_2^{2-} Chlorate
HPO_4^{2-} Hydrogen phosphate	O_2^{2-} Chlorite
$\text{C}_2\text{H}_3\text{O}_2^{-}$ Acetate	ClO^{-} Hypochlorite
OH^{-} Hydroxide	CrO_4^{2-} Chromate
CN^{-} Cyanide	$\text{C}_2\text{O}_4^{2-}$ Dioxalate
O_2^{2-} Peroxide	MnO_4^{-} Permanganate

Source: *Chemistry: Principles and Practice*, 6e, 2008, by T. L. Brown et al. Copyright 2008



Ionic Bonds and Compounds


Monatomic atoms

H ⁺																							
Li ⁺	Mg ²⁺																						
Na ⁺	K ⁺																						
K ⁺	Ca ²⁺																						
Rb ⁺	Sr ²⁺																						
		$\begin{matrix} \text{[OH}^-\text{]} \text{ [Mn}^{2+}\text{]} \text{ [Fe}^{2+}\text{]} \text{ [Co}^{2+}\text{]} \\ \text{[O}^{2-}\text{]} \text{ [Mn}^{3+}\text{]} \text{ [Fe}^{3+}\text{]} \text{ [Co}^{3+}\text{]} \end{matrix}$														$\begin{matrix} \text{Sc}^{3+} \\ \text{Cu}^{2+} \end{matrix}$	Zn ²⁺	$\begin{matrix} \text{Al}^{3+} \\ \text{Sn}^{2+} \\ \text{Pb}^{2+} \end{matrix}$	$\begin{matrix} \text{M}^{3+} \\ \text{O}^{2-} \end{matrix}$	$\begin{matrix} \text{P}^{3-} \\ \text{S}^{2-} \end{matrix}$	Cl ⁻	Br ⁻	I ⁻

Polyatomic atoms

NH ₄ ⁺ Ammonium																	
NO ₃ ⁻	Nitrate	SO ₄ ²⁻	Sulfate														
CO ₃ ²⁻	Carbonate	SO ₃ ²⁻	Sulfite														
HCO ₃ ⁻	Bicarbonate	HSO ₄ ⁻	Bisulfate														
	(Hydrogen carbonate)		(Hydrogen sulfate)														
NO ₂ ⁻	Nitrite	ClO ₄ ⁻	Perchlorate														
PO ₄ ³⁻	Phosphate	ClO ₃ ⁻	Chlorate														
H ₂ PO ₄ ⁻	Hydrogen phosphate	ClO ₂ ⁻	Chlorite														
C ₂ H ₃ O ₂ ⁻	Acetate	ClO ⁻	Hypochlorite														
OH ⁻	Hydroxide	CrO ₄ ²⁻	Chromate														
CN ⁻	Cyanide	Cr ₂ O ₇ ²⁻	Dichromate														
O ₂ ²⁻	Peroxide	MnO ₄ ⁻	Permanganate														

Source: Chemistry Online, Dec 19, 2003 by G. Martinez and C. Martinez




Ionic Bonds and Compounds, Continued



ionic bond – an attraction between oppositely charged ions


ionic compound – composed of charged ions

Metal cations and nonmetal anions form ionic compounds.



Ionic Compound Structure

ionic lattice – an array of positive and negative ions.



a

b

Book: Introductory Chemistry, 2e, © 2011 W. H. Freeman and Company




Figure 1 consists of two parts, (a) and (b). Part (a) shows a single repeating unit of the polymer, which is a central carbon atom (black) bonded to two nitrogen atoms (green) and two oxygen atoms (purple). The nitrogen atoms are labeled with a '+' sign, and the oxygen atoms are labeled with a '-' sign. Part (b) shows a long chain of repeating units, represented by a series of green and purple spheres connected by lines, forming a zigzag pattern.

Chemical Formulas

Show the type and amount of each element present


Empirical formula: The smallest whole-number ratio of atoms

Formula unit: The smallest number of ions necessary to form a compound




cation
 NaCl
anion

Formula unit: The smallest number of ions necessary to form a compound



Ionic Compounds

Write the formula for a compound composed of potassium and sulfide ions.



Total charge = 0




Revel, Introductory Chemistry, 2e © 2021 W. H. Freeman and Company


positive charges must equal the negative charges.

Compounds with Polyatomic Ions

Write the formula for a compound composed of calcium and nitrate ions.



Total charge = 0





Revel, Introductory Chemistry, 2e © 2021 W. H. Freeman and Company

positive charges must equal the negative charges.

Compounds with Polyatomic Ions, Continued

Write the formula for a compound composed of aluminum and sulfate ions.



positive charges must equal the negative charges.

Naming Ionic Compounds, Part 1

cation anion

NaCl *sodium chloride*

MgCl₂ *magnesium chloride*

MgSO₄ *magnesium sulfate*

Naming Ionic Compounds, Part 2


cation anion

Cu⁺
CuCl *copper(I) chloride*

Cu²⁺
CuCl₂ *copper(II) chloride*

Example, Naming Ionic Compounds

1. Name the compound Fe(NO₂)₂.




2. Write the empirical formula for ammonium sulfide.




Summary, Ionic Compounds

- Ionic bonds occur between oppositely charged ions
- In ionic compounds, total charge = 0
- Named as “cation anion”
- Formula $\leftrightarrow$ Name




Covalent Bonding, Part 1

covalent bond – electrons shared between two atoms




Covalent Bonding, Part 2

covalent bond – electrons shared between two atoms



By sharing electrons, each hydrogen completes its valence level.

Covalent Bonding, Part 3



Seven Elements Form Diatomic Molecules

The Magnificent Seven


Hydrogen: H_2
Nitrogen: N_2
Oxygen: O_2
Fluorine: F_2
Chlorine: Cl_2
Bromine: Br_2
Iodine: I_2

Revell, *Introductory Chemistry*, 2e, © 2021 W. H. Freeman and Company

Seven Elements Form Diatomic Molecules. Continued


[illegible]

Double and Triple Bonds in Lewis Structures



Covalent Compounds


Covalent compounds fulfill the octet rule by sharing electrons.



Revel, Introductory Chemistry, 2e, © 2021
W. H. Freeman and Company

Electrons in Lewis Structures

In this structure, how many electrons does the nitrogen atom share through covalent bonds? How many of the valence nitrogen electrons are not shared? Does this nitrogen atom have a complete octet?




Nitrogen has 6 shared electrons
 and 2 unshared electrons

 8 electrons - a complete octet


Covalent Compounds, Continued

molecular formula – gives the number of atoms in the molecule



Empirical Formula: NH_2
 Molecular Formula: N_2H_4

Covalent Compound Structures




Covalent compounds often have complex structures.

Compound name	Formula
Phosphorus monoxide	PO
Diphosphorus trioxide	P_2O_3
Diphosphorus tetroxide	P_2O_4
Tetraphosphorus decoxide	P_4O_{10}

Naming Binary Covalent Compounds

Atoms	Prefix
1	mono-
2	di-
3	tri-
4	tetra-
5	penta-
6	hexa-
7	hepta-
8	octa-
9	nona-
10	deca-

PCl_3 phosphorus trichloride
 PCl_5 phosphorus pentachloride



* omit for first element

Using Greek Prefixes

“pent” or “penta”

PCl_5 phosphorus **pent**achloride

P_2O_5 diphosphorus **pent**oxide

Remove “a” if anion begins with a vowel.

Practice Naming Covalent Compounds


Nitrogen and oxygen form two covalent compounds, NO_2 and N_2O_4 . Name each of these compounds.

NO_2 nitrogen **d**ioxide

N_2O_4 **d**initrogen **tet**roxide

Summary of Covalent Compounds

- In covalent bonds, atoms share electrons
- Covalent bonds form between nonmetals
- Most covalent compounds form discrete molecules
- We describe molecules using
 - Lewis structures
 - Molecular formulas
- Naming binary covalent compounds
 - Leftmost element first
 - Second element named as anion
 - Prefixes indicate the number of atoms present



Distinguishing Ionic and Covalent Compounds

- To fulfill the octet rule, atoms
- gain or lose electrons (ions)
 - share electrons (covalent bonds)


Covalent compounds

- share electrons
- between nonmetal atoms
- usually form molecules
- *molecular formula*



Ionic compounds

- oppositely-charged ions
- don't form molecules
- *formula unit* or *empirical formula*



Properties of Ionic and Covalent Compounds

Limestone
(CaCO_3)

Olive Oil



Identifying and Naming Compounds

Covalent compounds

- all nonmetals

Ionic compounds

- metal + nonmetal
- contains polyatomic ions

Identify these compounds as ionic or covalent, and name each one:

MgF_2
ionic
magnesium fluoride

$\text{Fe}(\text{NO}_3)_3$
ionic
iron(III) nitrate

P_2O_4
covalent
diphosphorus tetroxide

SCl_6
covalent
sulfur hexachloride

Aqueous Solutions:


How Ionic and Covalent Compounds Differ

aqueous solution A homogeneous mixture, in which the main component is water


soluble Able to dissolve



Electrolyte Solutions Conduct Electricity




dissociation Ions are pulled apart in an aqueous solution



Acids

covalent compounds that produce H^+ ions in aqueous solution



Common Acids

Formula	Name	Formula	Name
HF	hydrofluoric acid	HNO ₃	nitric acid
HCl	hydrochloric acid	HNO ₂	nitrous acid
HBr	hydrobromic acid	H ₂ SO ₄	sulfuric acid
HI	hydroiodic acid	H ₃ PO ₄	phosphoric acid
H ₂ CO ₃	carbonic acid	HC ₂ H ₃ O ₂	acetic acid

Binary Acids

HF	hydrofluoric acid
HCl	hydrochloric acid
HBr	hydrobromic acid
HI	hydroiodic acid

Oxyacids

form H^+ and oxyanion

1. -ate $\rightarrow$ -ic acid

NO ₃ ⁻	nitrate	HNO ₃	nitric acid
CO ₃ ²⁻	carbonate	H ₂ CO ₃	carbonic acid
SO ₄ ²⁻	sulfate	H ₂ SO ₄	sulfuric acid
PO ₄ ³⁻	phosphate	H ₃ PO ₄	phosphoric acid

Oxyacids, Continued


form H^+ and oxyanion

2. *-ite* $\rightarrow$ *-ous acid*

NO_2^-	nitrite	HNO_2	nitrous acid
ClO_2^-	chlorite	HClO_2	chlorous acid

Summary, Electrolytes


electrolytes $\left\{ \begin{array}{l} \text{ionic compounds} \\ \text{acids (form } \text{H}^+ \text{ ions in water)} \end{array} \right.$



Introductory Chemistry
Chem 103

Chapter 6 – Chemical Reactions


Lecture Slides




Chemical Equations




Chemical Equations Show Ratios of Substances



Chemical Equations Show Ratios of Substances, Continued



The Ratios In a Chemical Reaction Are Constant



$\frac{2,000,000}{\text{molecules}}$
 $\frac{1,000,000}{\text{molecules}}$
 $\frac{2,000,000}{\text{molecules}}$

In a **balanced equation**, the number and type of each atom are the same on both sides of the arrow.

Properly balanced – smallest whole-number ratio

Balancing Equations

In a **balanced equation**, the number and type of each atom are the same on both sides of the arrow.



Practice Balancing Equations



1. Identify number and type on each side.
2. Add coefficients to balance atoms.
3. Do not change subscripts.

Practice Balancing Equations, Continued



Balance elemental forms last.

Strategies for Balancing Equations

balance polyatomic ions




nitrate: NO_3^-

hydroxide: OH^-

Strategies for Balancing Equations, Continued



use a fractional coefficient for diatomic molecules



need 7 oxygen atoms!



Equations with Phase Notations

phase notations: show phase or state of reaction components




TABLE 6.1 Phase Symbols

Symbol	Meaning
(s)	Solid
(l)	Liquid
(g)	Gas
(aq)	Aqueous solution (dissolved in water)

Aqueous Solutions

(aq) – indicates the substance is dissolved in water



Chemical Equations Can Show Changes of State



Classifying Reactions, Part 1



Classifying Reactions, Part 2



Classifying Reactions, Part 3

Decomposition:

One forms two or more





Single Displacement:

One element replaces another





Synthesis (Combination):

Two form one



Double Displacement:

Two ions replace each other



Reuell, Introductory Chemistry, 2e, © 2021
W. H. Freeman and Company



Decomposition Reactions



Decomposition:
One forms two or more





Synthesis Reactions



Synthesis (Combination):
Two form one




Single Displacement Reactions



Single Displacement:
One element replaces another

Double Displacement Reactions




The anions "swap" positions




Double Displacement
Two ions replace each other

Single and Double Displacement Reactions




Classifying Reactions Summary


Decomposition:
One forms two or more



Synthesis (Combination):
Two form one



Single Displacement:
One element replaces another




Double Displacement:
Two ions replace each other



Reactions between Metals and Nonmetals

Metal + Nonmetal $\rightarrow$ Ionic Compound





metal cation + nonmetal anion



oxidation – loss of electrons

reduction – gain of electrons

Reactions between Metals and Nonmetals Example 1



Reactions between Metals and Nonmetals Example 2



Mg loses two electrons

Metals and Nonmetals Form Specific, Stable Ions.

[illegible]

Inductron Chemistry 2e © 2021 W H Freeman and Company


Reactions Between Metals and Nonmetals Practice

What compound is formed when aluminum metal reacts with chlorine gas? Write a balanced equation for this reaction.






Reactions Between Metals and Nonmetals, More Practice

When tin metal reacts with bromine, it is oxidized to the tin(IV) ion, while bromine is reduced to form bromide ions. Write a balanced equation for this reaction.



Combustion Reactions

reactions in which oxygen gas combines with elements or compounds to produce oxides.



Hydrocarbons compounds composed of hydrogen and carbon



TABLE 6.2 Common Hydrocarbons

Formula	Name	Use
CH ₄	Methane	Natural gas
C ₂ H ₂	Acetylene	Torches for cutting and welding
C ₂ H ₄	Ethylene	Manufacture of plastic
C ₃ H ₈	Propane	Natural gas component; used for heating and power
C ₄ H ₁₀	Butane	Lighter fluid
C ₆ H ₆	Benzene	Solvent; precursor for many pharmaceutical compounds
C ₈ H ₁₈	Octane	Component of gasoline



Combustion of Hydrocarbons

hydrocarbon + oxygen → carbon dioxide + water



The Combustion of Sulfur Produces Sulfur Oxides


Sulfur Oxides
(SO_x)

- SO
- SO₂
- SO₃
- SO_x
- S₂O₇



Combustion Reactions Practice

Write a balanced equation for the combustion of calcium metal.




Combustion Reactions, More Practice

Write a balanced equation for the combustion of propane gas, a common fuel used for home heating, cooking, etc. The formula for propane is C₃H₈.



Reactions in Aqueous Solution

Ionic compounds **dissociate** when dissolved in water.



Comparing Molecular and Ionic Equations

molecular equation – shows ions together as compounds





ionic equation – shows dissociated ions as separate species




Writing Ionic Equations Practice

Show this process as an ionic equation:




Predicting Solubility

Many ionic compounds are **insoluble** in water.



Predicting Solubility, Continued



Factors affecting solubility

- Charge on ions
- Size of ions
- How tightly ions pack together

Soluble

NaCl (Na⁺ and Cl⁻)
KNO₃ (K⁺ and NO₃⁻)
NH₄Br (NH₄⁺ and Br⁻)

Insoluble

Fe₂O₃ (Fe³⁺ and O²⁻)
PbS (Pb²⁺ and S²⁻)
BaCO₃ (Ba²⁺ and CO₃²⁻)

Solubility Rules:

- Halogens (F^- , Br^- , Cl^- , I^-) are soluble
– Unless bonded to Ag^+ or Pb^{2+}

Soluble	Insoluble
KF	AgF
ZnCl ₂	AgCl
FeBr ₂	PbBr ₂
CuI	PbI ₂

Solubility Rules, Continued

TABLE 6.3 Solubility Rules

Compounds Containing These Ions Are Nearly Always Soluble	
Alkali metals	Li^+ , Na^+ , K^+ , Rb^+
Ammonium	NH_4^+
Large -1 oxyanions	NO_3^- , ClO_3^- , ClO_4^- , $C_2H_3O_2^-$
Compounds Containing These Ions Are Usually Soluble	
Halides (except Pb^{2+} , Ag^+)	F^- , Cl^- , Br^- , I^-
Sulfate (except Ba^{2+} , Ca^{2+} , Pb^{2+} , Ag^+)	SO_4^{2-}
Not Soluble	
Most other ions	

Solubility Tables

	Ammonium	Sodium	Potassium	Silver	Magnesium	Calcium	Bismuth	Zinc	Copper(I)	Copper(II)	Lead(II)	Iron(II)	Aluminum
Nitrate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Chlorate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Perchlorate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Acetate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Chromate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Bromide	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Iodide	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Sulfate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Sulfite	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Carbonate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Phosphate	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes

Ex.: $CaCl_2$ soluble
 $Mg(OH)_2$ insoluble

Determine Solubility

Determine whether the following compounds are soluble or insoluble in water:

Na_3PO_4 soluble $AlCl_3$ soluble $CaCO_3$ insoluble


TABLE 6.3 Solubility Rules

Compounds Containing These Ions Are Nearly Always Soluble	
Alkali metals	Li^+ , Na^+ , K^+ , Rb^+
Ammonium	NH_4^+
Large -1 oxyanions	NO_3^- , ClO_3^- , ClO_4^- , $C_2H_3O_2^-$
Compounds Containing These Ions Are Usually Soluble	
Halides (except Pb^{2+} , Ag^+)	F^- , Cl^- , Br^- , I^-
Sulfate (except Ba^{2+} , Ca^{2+} , Pb^{2+} , Ag^+)	SO_4^{2-}
Not Soluble	
Most other ions	



Precipitation Reactions

precipitation reaction two aqueous solutions produce an insoluble product

precipitate the solid product formed in the reaction




$Pb(NO_3)_2(aq)$




$NaI(aq)$


Precipitation Reactions Are Double Displacement Reactions




The anions "swap" positions



How Precipitation Reactions Occur




Driving force - formation of the solid




Comparing Complete and Net Ionic Equations

Complete ionic equation
shows all ions present



Net ionic equation
Only include ions involved in the precipitation



Writing Precipitation Reactions

Three ways to show a precipitation reaction:


Molecular Equation <i>shows neutral compounds</i> $\text{Pb}(\text{NO}_3)_2(\text{aq}) + 2 \text{KCl}(\text{aq}) \rightarrow \text{PbCl}_2(\text{s}) + 2 \text{KNO}_3(\text{aq})$
Complete Ionic Equation <i>shows all ions present</i> $\text{Pb}^{2+}(\text{aq}) + 2 \text{NO}_3^{-}(\text{aq}) + 2 \text{K}^{+}(\text{aq}) + 2 \text{Cl}^{-}(\text{aq}) \rightarrow \text{PbCl}_2(\text{s}) + 2 \text{K}^{+}(\text{aq}) + 2 \text{NO}_3^{-}(\text{aq})$
Net Ionic Equation <i>Omits spectator ions; only shows ions that react.</i> $\text{Pb}^{2+}(\text{aq}) + 2 \text{Cl}^{-}(\text{aq}) \rightarrow \text{PbCl}_2(\text{s})$

Use solubility rules to predict precipitation reactions.

Precipitation Reactions Practice



When aqueous silver acetate is combined with aqueous barium chloride, a white precipitate forms. Write balanced complete ionic, net ionic, and molecular equations to show the reaction that takes place. Include phase symbols.



Complete ionic equation



Net ionic equation



Molecular equation



Summary of Precipitation Reactions


- Soluble ionic compounds dissociate in water.
- Some ionic compounds are insoluble in water.
- Solubility rules predict the solubility of compounds.
- Precipitation reaction: two solutions combine to produce an insoluble product.
- We describe reactions in solution using
 - molecular equations
 - complete ionic equations
 - net ionic equations

Reactions in Aqueous Solution

acids compounds that produce H^{+} ions in aqueous solution

TABLE 6.4 Common Acids

Formula	Name
HF	Hydrofluoric acid
HCl	Hydrochloric acid
HBr	Hydrobromic acid
HI	Hydroiodic acid
H_2CO_3	Carbonic acid
HNO_3	Nitric acid
HNO_2	Nitrous acid
H_2SO_4	Sulfuric acid
H_3PO_4	Phosphoric acid
$\text{HC}_2\text{H}_3\text{O}_2$	Acetic acid



Reactions in Aqueous Solution, Continued

bases compounds that produce OH^- ions in aqueous solution





TABLE 6.5 Common Hydroxide Base



Formula	Name
LiOH	Lithium hydroxide
NaOH	Sodium hydroxide
KOH	Potassium hydroxide
Ba(OH)_2	Barium hydroxide

Neutralization Reactions


Acids and bases undergo **neutralization reactions**.



Ex.: hydrochloric acid reacts with sodium hydroxide





Ex.: nitric acid reacts with lithium hydroxide



Neutralization Reactions, Continued

Acid-base neutralization is a **double displacement reaction**.



The formation of water is the driving force for the reaction.

Acid-Base Reactions Practice

Write a balanced equation to show the reaction of sulfuric acid with sodium hydroxide. Include phase symbols.

